

CRITICAL CARE STABILISATION GUIDE

An aide-memoire to support the Obstetric Anaesthetic Team caring for a ventilated COVID-19 obstetric patient. To be completed with every patient review.

**For urgent concerns & referrals
contact ICU consultant bleep 5490**

**NO EXCEPTIONS: PPE
BEFORE ENTERING ROOM**

Patient addressograph

SEDATION

Propofol 1% (range 5-30ml/h) _____ ml/hr

Alfentanyl (500mcg/ml) (range 1-5ml/hr) _____ ml/hr

Is an epidural in situ? Yes / No Checked

LINES

Arterial line Site _____ Checked

IV line 1 Site _____ Checked

IV line 2 Site _____ Checked

CVC Site _____ Checked

Urinary Catheter Size _____ Checked

NGT Size _____ Secured @ _____ cm Checked

VENTILATOR

ETT Size _____ Secured @ _____ cm Checked

FiO2 (Target SpO2>94%) = _____ %

TV (Target 6ml/kg) = _____ ml

PEEP (Initially 5-10cmH2O) = _____ cmH₂O

CXR - confirm ETT position Checked

- confirm NGT position Checked

- confirm CVC position Checked

CARDIOVASCULAR

MAP (Target >65mmHg) = _____ mmHg

Noradrenaline (Initially 4mg in 50ml) _____ ml/hr

Fluid balance (minimise to maintain oxygenation) +/- _____ ml/24h

OBSTETRIC COMPLICATION(s) management plan reviewed -

Completed by: _____ (Please print) Date: _____ Time: __:__ Location _____

